



## Identification of Active Fatty Acid Mixture Extracted from Rice Bran Oil Using NMR Spectroscopy

Patamapan Tamvapee<sup>1</sup>, Siritron Samosorn<sup>2</sup>, Ramida Watanapokasin<sup>1\*</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Medicine, Srinakharinwirot University, Bangkok, Thailand

<sup>2</sup>Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Srinakharinwirot University, Bangkok, Thailand

### ARTICLE INFO

#### Article history:

Received 01 April 2023

Revised 23 April 2023

Accepted 25 April 2023

Published online 01 May 2023

**Copyright:** © 2023 Tamvapee *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### ABSTRACT

Rice bran oil (RBO) is oil obtained from rice bran. It contains fatty acids and some phytochemicals. RBO has been reported to have some biological activities. Previous publications showed the anticancer activity of rice bran oil fraction extracted from rice bran oil in LoVo, a colorectal cancer cell line through apoptosis via the MAPK pathway with the half-maximal effective concentration (EC<sub>50</sub>) value of 155.50 ± 1.10 µg/ml. Thus, this study investigated the fatty acid composition of the active rice bran oil fraction purified by column chromatography and identified by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy. The result showed that the active fatty acid fraction consists of saturated fatty acid and unsaturated fatty acid in a ratio of 36:64. These findings may lead to the development of alternative cancer therapies in the future.

**Keywords:** Rice bran oil; fatty acid; NMR spectroscopy

### Introduction

Cancer is a growing threat to human health worldwide, in some cases, may involve mass morbidity and mortality.<sup>1</sup> Natural products such as rice bran oil offer biological benefits and have inspired the development of new medicines.<sup>2</sup>

Rice (*Oryza sativa L.*) is a vital staple worldwide.<sup>3</sup> Rice bran (RB) is a by-product of the rice grain milling process. It accounts for about 10% of the total weight of whole rice grain and is mainly comprised of aleurone, subaleurone layer, pericarp, and germ. Rice bran oil (RBO) extracts from RB's lipid part, and the specific content varies between 10-23%. RBO generally comprises unsaturated fatty acids (55-87%) such as oleic acid and linoleic acid, and saturated fatty acids (19-35%) such as palmitic acids.<sup>4</sup> RBO also contains functional components (2-3%) such as tocopherol (vitamin E), tocotrienol,  $\gamma$ -oryzanol, and phytosterol. RBO has been demonstrated to have cholesterol-lowering, antihypertension, hyperglycemia-lowering,<sup>2</sup> antidiabetic, antioxidant, and anticancer effects.<sup>5</sup>

In addition to rice bran oil, several studies showed the anticancer activity of natural compounds or oils that contain fatty acids. For example, Apple seed oil contained oleic acid (47%), linoleic acid (44%), palmitic (7%), stearic (2%), and arachidic acids (1%) showed good pharmacological potential against A549, a human lung carcinoma and SiHa, a human cervical cancer cell line.<sup>6</sup> Almond oil from Northern Cyprus and Turkey contained oleic acid (78%; 75%), linoleic acid (14%; 16%), and palmitic acid (7%; 6%), both exhibited significant anti-proliferative and anticancer activity in Colo-320 and Colo-741, human colon adenocarcinoma cell lines.<sup>7</sup>

\*Corresponding author. E mail: [ramidaw@g.swu.ac.th](mailto:ramidaw@g.swu.ac.th)  
[ramidabc@gmail.com](mailto:ramidabc@gmail.com)  
Tel: +66824797824

**Citation:** Tamvapee P, Samosorn S, Watanapokasin R. Identification of Active Fatty Acid Mixture Extracted from Rice Bran Oil Using NMR Spectroscopy. Trop J Nat Prod Res. 2023; 7(4) <http://www.doi.org/10.26538/tjnpr/v7i4.14>

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

The essential oil extracted from *Pinus roxburghii* showed anticancer activity in many human cell lines, especially HCT-116, a human colorectal carcinoma, and KBM-5, a human myelogenous leukemia cell lines.<sup>8</sup> Moreover, Essential Oil extracted from *Arnica montana L.* showed anticancer activity by induced apoptosis in MGGCCM, a human anaplastic astrocytoma, and T98G, a human glioblastoma multiforme cell lines.<sup>9</sup> Different types of coconut oil showed different anticancer efficiency in cell lines due to their fatty acid composition.<sup>10</sup> Oil extracted from *Zanthoxylum bungeanum* seed which has unsaturated fatty acid as a main component, showed anticancer activity by induction of apoptosis via MAPK pathway on A375, a human malignant melanoma cell line.<sup>11</sup> Besides direct anticancer activity, fatty acids are also used to produce fatty acid-conjugated therapeutic molecules.<sup>12, 13</sup> For example, the mixture of linoleic acid (CLA) reduced key enzymes in glycogenesis expression.<sup>14</sup> Other research found CLA have immunomodulatory effects on some functions of bovine monocytes.<sup>15</sup> Docetaxel, breast cancer therapeutic drug conjugated linoleic acid (DTX-LA) increased drug delivery efficiency and significant antitumor activity.<sup>16</sup> Fatty acids also show other biological activities, for example, Chloroform extraction of *Chrysophyllum albidum* leaf and stem-bark have antioxidant property.<sup>17</sup> Similar to essential oils extracted from *Haplophyllum tuberculatum*<sup>18</sup> and *Dennettia tripetala*.<sup>19</sup> Some oils, such as coconut and moringa, showed protection against cadmium-induced toxicity in rats.<sup>19</sup>

Nuclear magnetic resonance (NMR) is a tool used to characterize organic compounds' structures. It is a very reliable technique and does not require chemical standards.<sup>20</sup> NMR is also beneficial in providing structural information and quantifying chemical composition in mixture samples. Quantification can be determined by peak area per nuclei, which is proportional to the number of corresponding nuclei. Several precedents report quantifying fatty acid composition in vegetable oils by proton (<sup>1</sup>H) NMR.<sup>21-24</sup> Integration values of relevant proton signals were used to determine the amounts of fatty acids in vegetable oils.

Previous study showed anticancer activity of fatty acid fraction extracted from rice bran oil.<sup>25</sup> The fraction induced apoptosis in LoVo, colorectal cancer cell line through MAPK pathway. Thus, in this study, we aim to determine the ratio of saturated and unsaturated fatty acids of the active RBO fraction from integration values of characteristic proton signals of each fatty acid.

## Materials and Methods

### Materials

Organic rice bran oil extract (RBOE) Lot. 27062231 (June 22, 2017) was obtained from the Bakrua farmer group in Yasothon province, Thailand. All the solvents were analytical reagent grade or distilled. Column chromatography was performed under gravity on Merck Kieselgel 60 silica gel (230-400 mesh). All fractions were monitored by thin-layer chromatography (Merck Silica gel 60 F<sub>254</sub> with a thickness of 0.2 mm on an aluminum sheet). The separated bands were detected by UV lamp at 254 nm and 366 nm and visualization with iodine vapours. NMR spectra were recorded on a Bruker AVANCE III 400 NMR spectrometer operating at 400 MHz for <sup>1</sup>H and 100 MHz for <sup>13</sup>C in deuterated chloroform using TMS as an internal standard.

### Isolation and Characterization

The crude RBOE (62.63 g) was added with hexane (600 ml) and partitioned with 90% aqueous methanol (600 ml) three times. The combined aqueous methanol layers were concentrated in a vacuum at 40°C to give 6.85 g. The extract was subjected to silica column chromatography with chloroform-methanol with gradually increasing methanol concentrations to give 11 fractions (FAs). All fractions were monitored by thin-layer chromatography (TLC) and screening for cancer cell viability using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) assay. The FA3 (0.32 g) was eluted with 2-3% methanol in chloroform. It showed two spots on TLC, which was further purified by a silica column chromatography with 2% methanol in chloroform to afford an off-white soft wax in oil (0.04 g) for anticancer activity testing and chemical composition profile by NMR spectroscopic technique.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 5.32-5.35 (m, 6H, -CH=CH-), 2.75 (t, *J* = 4.0 Hz, 2H, =CHCH<sub>2</sub>CH=), 2.33 (t, *J* = 8.0 Hz, 20H, CH<sub>2</sub>COOH), 1.99-2.03 (m, 26H, CH<sub>2</sub>CH=CH-), 1.61 (t, *J* = 8.0 Hz, 29H, CH<sub>2</sub>CH<sub>2</sub>COOH), 1.20-1.50 (m, 225H, -(CH<sub>2</sub>)<sub>n</sub>), 0.86 (t, *J* = 4.0 Hz, 38H, -CH<sub>3</sub>).

<sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 14.1, 22.6, 22.7, 24.7 (2C), 25.7, 27.2 (3C), 29.0 (2C), 29.2, 29.3(2C), 29.4, 29.5, 29.6, 29.7, 29.8, 31.6, 31.9 (2C), 33.9, 127.9, 128.1, 129.8, 130.0, 130.2, 179.2.


## Results and Discussion

MTT results from previous research showed that rice bran oil extract fraction (FA3) has the best effect on LoVo cells viability with the half-


maximal effective concentration (EC<sub>50</sub>) value of 155.50 ± 1.10 µg/ml. In contrast, it showed no inhibition against the control cells at the same concentration, Vero, with an EC<sub>50</sub> of 258.60 ± 1.07 µg/ml. Then this fraction was used to determine the molecular mechanism against LoVo cells, and found that it able to induced apoptosis in LoVo cells through the MAPK pathway.<sup>25</sup>

Isolation of the FA3 fraction was obtained from silica column chromatography as an off-white oil wax showing only one spot on TLC with 1% methanol in dichloromethane. The nuclear magnetic resonance (NMR) spectroscopic data of the FA3 indicated that it contained a mixture of saturated and unsaturated fatty acids. Typically, fatty acids obtained from rice bran oil extract contained fatty acids (Figure 1), including palmitic acid as a major saturated fatty acid, oleic acid as a major monounsaturated fatty acid, and linoleic acid as a major polyunsaturated fatty acid.<sup>26, 27</sup>

The FA3 NMR data were identified by comparing the obtained <sup>1</sup>H and <sup>13</sup>C NMR spectra with those of previously reported spectra. The <sup>1</sup>H-NMR spectrum (Figure 2) had no signals for methylene and methine protons in α-position of glycerol backbone in the range of 4.0-4.5 ppm, which indicated that triglyceride was not present in this fraction. The spectrum showed signals assigned to olefinic protons of unsaturated fatty acids at 5.32-5.35 ppm. The signal at 2.75 ppm was assigned to the *bis*-allylic proton of polyunsaturated fatty acid, and the signal at 1.99-2.03 ppm was assigned to allylic protons of unsaturated fatty acid. The signals at 2.33 and 1.61 ppm were ascribed to α- and β-methylene protons adjacent to the carbonyl group, respectively. The signals assigned to methylene protons on saturated carbon atoms and terminal methyl protons were observed at 1.20-1.50 and 0.86 ppm, respectively.




**Figure 1:** Major fatty acids composition of rice bran oil. [18]



**Figure 2:** <sup>1</sup>H NMR spectrum of the FA3 with integration values.

The  $^{13}\text{C}$  NMR spectrum showed a downfield signal attributed to the carbonyl carbon of a carboxylic acid at 179.2 ppm. The characteristic signals were ascribed to olefinic carbon of unsaturated fatty acid in the 127-131 ppm region, as shown in Figure 3.<sup>28</sup>

Each peak area in the  $^1\text{H}$ -NMR spectrum is proportional to the number of protons of each signal. Therefore, the content of saturated and unsaturated fatty acids can be obtained by measuring the peak area of each fatty acid characteristic signal.  $\alpha$ -methylene proton adjacent to the carbonyl group (signal A) and allylic proton (signal B) in Figure 2 is helpful for this purpose.<sup>29</sup> Therefore, the total amounts of saturated and unsaturated fatty acids can be determined by using signal A at 2.33 ppm, and that of unsaturated fatty acids can be determined by signal B at 1.99-2.03 ppm. Subsequently, the content of saturated fatty acids can also be determined by assuming that the total content (C) of fatty acids equals 100% and then subtracting the content of unsaturated fatty acids ( $\%C_{\text{sat}} = 100 - C_{\text{unsat}}$ ). The integration value ratio of signal A of total fatty acids to signal B of unsaturated fatty acid is two  $\alpha$ -methylene protons (integration value/2:  $20.17/2 = 10.09$ ) to two four allylic protons (integration value/4:  $25.93/4 = 6.48$ ). Then converting this integration value ratio into percentages and found a ratio of 64.2% unsaturated fatty acid to 35.8% saturated fatty acid is obtained.



**Figure 3:**  $^{13}\text{C}$  NMR spectrum of the FA3.

### Conclusion

In the present study, NMR analysis of FA3 demonstrated that it contained amounts of saturated fatty acid: unsaturated fatty acid in the ratio of 36:64. This oil mixture exhibited anti-cancer activity as in the previous report.

### Conflict of Interest

The authors declare no conflict of interest.

### Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

### Acknowledgments

Special appreciation to the Bakrua farmers group in Yasothon province, Thailand, for their support and for providing access to the resources necessary to complete this study. We appreciate, Professor Hitomi Ohara and Dr. Wichean Khawdas, Faculty of Fiber Science and Engineering, Kyoto Institute of Technology, Japan, for their kind assistance recording NMR data. Our gratitude to the Research Division,

Faculty of Medicine, and Department of Chemistry, Srinakharinwirot University, for providing equipment to conduct experiments.

### Funding

This work was supported by the Research and Researcher for Industries (RRI), Thailand Research Fund under Grant PHD5710023.

### References

1. Ma X, Yu H. Global burden of cancer. *Yale J Biol Med.* 2006;79(3-4):85-94.
2. Wang Y. Chapter 6 - Applications of Rice Bran Oil. In: Cheong L-Z, Xu X, editors. *Rice Bran and Rice Bran Oil*: AOCS Press; 2019. p. 159-68.
3. Rigo LA, Pohlmann AR, Guterres SS, Ruver Beck RC. Chapter 23 - Rice Bran Oil: Benefits to Health and Applications in Pharmaceutical Formulations. In: Watson RR, Preedy VR, Zibadi S, editors. *Wheat and Rice in Disease Prevention and Health*. San Diego: Academic Press; 2014. p. 311-22.
4. Dunford NT. Chapter 1 - Chemistry of Rice Bran Oil. In: Cheong L-Z, Xu X, editors. *Rice Bran and Rice Bran Oil*: AOCS Press; 2019. p. 1-18.
5. Lai O-M, Jacoby JJ, Leong W-F, Lai W-T. Chapter 2 - Nutritional Studies of Rice Bran Oil. In: Cheong L-Z, Xu X, editors. *Rice Bran and Rice Bran Oil*: AOCS Press; 2019. p. 19-54.
6. Walia M, Rawat K, Bhushan S, Padwad YS, Singh B. Fatty acid composition, physicochemical properties, antioxidant and cytotoxic activity of apple seed oil obtained from apple pomace. *J Sci Food Agric.* 2014;94(5):929-34.
7. Mericli F, Becer E, Kabadayi H, Hanoglu A, Yigit Hanoglu D, Ozkum Yavuz D, Ozek T, Vatansever S. Fatty acid composition and anticancer activity in colon carcinoma cell lines of *Prunus dulcis* seed oil. *Pharm Biol.* 2017;55(1):1239-48.
8. Sajid A, Manzoor Q, Iqbal M, Tyagi AK, Sarfraz RA, Sajid A. *Pinus Roxburghii* essential oil anticancer activity and chemical composition evaluation. *Excli j.* 2018;17:233-45.
9. Sugier D, Sugier P, Jakubowicz-Gil J, Winiarczyk K, Kowalski R. Essential Oil from *Arnica Montana* L. Achenes: Chemical Characteristics and Anticancer Activity. *Molecules.* 2019;24(22).
10. Verma P, Naik S, Nanda P, Banerjee S, Naik S, Ghosh A. In Vitro Anticancer Activity of Virgin Coconut Oil and its Fractions in Liver and Oral Cancer Cells. *Anticancer Agents Med Chem.* 2019;19(18):2223-30.
11. Pang W, Liu S, He F, Li X, Saira B, Zheng T, Chen J, Dong K, Pei XF. Anticancer activities of *Zanthoxylum bungeanum* seed oil on malignant melanoma. *J Ethnopharmacol.* 2019;229:180-9.
12. Bhat M, Jatyan R, Mittal A, Mahato RI, Chitkara D. Opportunities and challenges of fatty acid conjugated therapeutics. *Chem Phys Lipids.* 2021;236:105053.
13. Jóźwiak M, Filipowska A, Fiorino F, Struga M. Anticancer activities of fatty acids and their heterocyclic derivatives. *Eur J Pharmacol.* 2020;871:172937.
14. Szykowska A, Siwiec E, Gutowska I, Baranowska-Bosiacka I, Szczuko M, Kotlega D, Marlicz W, Stachowska E. Conjugated linoleic acids diminish glycogen synthase and glycogen synthase kinase-3 expression in muscle cells of C57BL/6J mice - *in vitro* and *in vivo* study. *J Physiol Pharmacol.* 2019;70(6).
15. Ávila G, Catozzi C, Pravettoni D, Sala G, Martino P, Meroni G, Lecchi C, Cecilian F. *In vitro* effects of conjugated linoleic acid (CLA) on inflammatory functions of bovine monocytes. *J Dairy Sci.* 2020;103(9):8554-63.
16. Zhang T, Li M, Yang R, Zhang D, Guan J, Yu J, Yang B, Zhang H, Zhang S, Liu D, Wang Y. Therapeutic efficacy of lipid emulsions of docetaxel-linoleic acid conjugate in breast cancer. *Int J Pharm.* 2018;546(1-2):61-9.

17. Ezema O, Omeje BO, Ozioko KN, Urama JC, Omeje DC, Nnawulezi H, Ejim A. A. Cholinesterase Inhibition, Biological Activity and Characterization of *Chrysophyllum albidum* Leaf and Stem-Bark Chloroform Extract Using GC-MS: An *In Vitro* Study. Trop J Nat Prod Res. 2021;5(1):130-4.
18. Agour A, Mssillou I, Saghrouchni H, Bari A, Lyoussi B, Derwich E. Chemical Composition, Antioxidant Potential and Antimicrobial Properties of the Essential Oils of *Haplophyllum tuberculatum* (Forsskal) A. Juss from Morocco. Trop J Nat Prod Res. 2020;4(12):1108-15.
19. Adedayo BC, Adebayo AA, Oboh G. *In Vitro* Antioxidant and Anti-Cholinesterase Properties of Essential Oils from Pepper Fruits (*Dennettia Tripetala* G. Baker). Trop J Nat Prod Res. 2020;4(9):596-600.
20. Teipel JC, Hausler T, Sommerfeld K, Scharinger A, Walch SG, Lachenmeier DW, Kuballa T. Application of <sup>1</sup>H Nuclear Magnetic Resonance Spectroscopy as Spirit Drinks Screener for Quality and Authenticity Control. Foods. 2020;9(10).
21. Matsui T, Iwasaki H, Matsumoto K, Osajima Y. Quality Evaluation of Edible Oils by Proton Nuclear Magnetic Resonance Measurement. Food Sci Technol. 1995;1(2):94-7.
22. Miyake Y, Yokomizo K, Matsuzaki N. Determination of unsaturated fatty acid composition by high-resolution nuclear magnetic resonance spectroscopy. J Am Oil Chem Soc. 1998;75(9):1091-4.
23. Guillen M, Ruiz A. <sup>1</sup>H Nuclear Magnetic Resonance as a fast tool for determining the proportions of the different acyl groups in acylglycerol mixtures. Eur J Lipid Sci Technol. 2003;105:502-7.
24. Barison A, da Silva CW, Campos FR, Simonelli F, Lenz CA, Ferreira AG. A simple methodology for the determination of fatty acid composition in edible oils through <sup>1</sup>H NMR spectroscopy. Magn Reson Chem. 2010;48(8):642-50.
25. Tamvapee P, Watanapokasin R. Apoptosis Induction through MAPK Signaling Pathway in LoVo Cells by Fatty Acid Fraction from Rice Bran Oil. Nutr Cancer. 2022;74(6):2122-32.
26. Olabanji I, Ajayi O, Adekunle A. Fatty acids, metal composition and physico-chemical parameters of Igbemo Ekiti rice bran oil. J Environ Chem Ecotoxicol. 2013;Volume 5:Page 39-46.
27. Latha RB, Nasirullah DR. Physico-chemical changes in rice bran oil during heating at frying temperature. J Food Sci Technol. 2014;51(2):335-40.
28. Miyake Y, Yokomizo K, Matsuzaki N. Rapid determination of iodine value by <sup>1</sup>H nuclear magnetic resonance spectroscopy. J Am Oil Chem Soc. 1998;75(1):15-9.
29. Guillén MaD, Ruiz A. High resolution <sup>1</sup>H nuclear magnetic resonance in the study of edible oils and fats. Trends Food Sci Technol. 2001;12(9):328-38.